

obtained a second peak, that indicates the plasma electrons. We integrated the plasma signal for 50P state, which shows the plasma lifetime (inset Fig 1 (a)). The integrated plasma signals for all nP states are shown in the Fig 1 (b). The study of the dynamics of spontaneous ionization as a function of delay time for a broad range of initial quantum states serves to characterize the n -dependence of Rydberg–Rydberg collisional ℓ -mixing, Penning ionization, and electron impact avalanche to plasma^[2,3]. In Fig. 1 (b) the higher states correspond to hot collisions and blackbody radiation, the middle states indicate the dipole-dipole interaction, ℓ -mixing, and n -mixing, while the lower states show the Penning ionization.

References

- [1] M. P. Robinson, B. L. Tolra, M. W. Noel, et al., Phys. Rev. Lett., 85 (2000)4466.
- [2] A Walz-Flannigan, J. R. Guest, JH Choi, et al., Phys. Rev. A, 69(2004)063405.
- [3] J. Hung, H. Sadeghi, M. Schulz-Weiling, et al., J. Phys. B: At. Mol. Opt. Phys., 47(2014)155301.

* Foundation item: National Natural Science Foundation of China (11274316, 21203216, 11404346)

3 - 11 Dielectronic Recombination of $^{56}\text{Fe}^{17+}$ at the CSRm*

Khan Nadir¹, Huang Zhongkui¹, Wen Weiqiang¹, Xu Xin², Wang Shuxing², Wang Hanbing¹, Dou Lijun¹, Zhu Xiaolong¹, Mao Lijun¹, Li Jie¹, Ma Xiaoming¹, Yang Jiancheng¹, Yuan Youjin¹, Zhu Linfan² and Ma Xinwen¹

(¹Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China;

²Department of Modern Physics, University of Science and Technology of China, Hefei Anhui 230026, China)

The ionization and heating of the media surrounding accretion-powered compact sources, such as cataclysmic




Fig. 1 (color online) The experimental DR rate coefficients for F-like $^{56}\text{Fe}^{17+}$ ions versus the collisional energy between electron and the ion beam in the c.m. frame. The DR resonances involve $\Delta n = 0$ transition. The observed DR channels are $\text{Fe}^{17+} (2s^2 2p^5 [^2P_{3/2}]) + e^- \rightarrow \text{Fe}^{16+} (2s^2 2p^5 [^2P_{1/2}] nl) (n = 18, \dots)$ and $\text{Fe}^{17+} (2s^2 2p^5 [^2P_{3/2}]) + e^- \rightarrow \text{Fe}^{16+} (2s^2 2p^6 [^2S_{1/2}] nl) (n = 6, \dots)$ respectively, which are indicated by vertical bars.

variables, X-ray binaries, and active galactic nuclei, are dominated by photoionization. Theoretical models of photo ionized gases show that the ionization structure is determined by photoionization balanced by radiative recombination (RR) and dielectronic recombination (DR). At low electron temperature existing in photo ionized gases with cosmic abundances, DR proceeds primarily via $nl_j \rightarrow nl'_j$, excitations of core electrons ($\Delta n = 0$). While Iron is the most abundant heavy element in astrophysical studies^[1]. So to address the needs for modeling photo ionized gases, we have initiated an experiment to measure the $\Delta n = 0$ DR rate for iron L-shell ions. Measurement were carried out by employing electron-ion merged-beam technique at heavy ion storage ring CSRm at IMP^[2]. Here we report our results for $\Delta n = 0$ DR of Fe^{17+} , where two series of resonance peaks of $^2P_{3/2} - ^2P_{1/2}$ and $^2P_{3/2} - ^2S_{1/2}$ (for excitation energies 12.718 2 eV and 132.006 2 eV respectively) were observed. However because of the instability of beam line and the breakdown of electron voltage detuning system, the series limit of DR channel $^2P_{3/2} - ^2S_{1/2}$ was not observed in these measurement as shown in Fig. 1. The further data analysis and the theoretical calculations are in progress.

References

- [1] D. W. Savin, T. Bartsch, M. H. Chen, et al., The Astrophysical Journal, 489(1997)L115.
- [2] Z K Huang, W Q Wen, H B Wang, et al., Phys. Scr. T166(2015)014023.

* Foundation item: National Natural Science foundation of China (11320101003) and External Cooperation Program of the CAS, (GJHZ1305)