

3 - 7 Observation of Indirect (e, 3e) of CO Induced by Electron Impact*

Zhang Pengju¹, Yan Shuncheng¹, Shen Lili^{1,2} and Ma Xinwen¹

(¹The Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China;

²University of Chinese Academy of Sciences, Beijing 100049, China)

We report an experimental investigation of ionization dissociation of CO^{2+} induced by electron impact at incident energy of 380 eV. The fragment ions (C^+/O^+) and the emitted electron were measured in coincidence by utilizing COLTRIMS technique. The kinetic-energy-release (KER) of the fragment ions and the energy of the outgoing electrons are obtained. Based on the characters of the KER distribution and the energy spectrum of the emitted




Fig. 1 (color online) Two-dimensional diagram of the kinetic energy of one of the emitted electrons (vertical) versus the KER of the ionic fragments (horizontal).

electrons (see Fig. 1), the direct and indirect double ionization ((e, 3e)) including an auto-ionization process were identified respectively.

Two apparent features are found as marked in Fig. 1. In the area A (marked by the red square), most of the emitted electrons distribute in the region with the electron kinetic energy less than 2 eV and the KER from 0 to 5 eV. In the area B (marked by the black square), the electrons distribute in the energy range of 0 to 14 eV and the intensity shows a monotonous decrease with the electron energy increase, and most of the electrons locate below 6 eV. The corresponding KER exhibits a broader distribution from 5 to 12 eV with a maximum around 7 eV. For the area A, the lower KER and the concentrated energy distribution of the emitted electron reveal the indirect (e, 3e) including an auto-ionization process^[1]. For the area B, the higher KER value and the continuous energy distribution of the outgoing electron indicate the direct (e, 3e) process.

Reference

- [1] T. Osipov, T. Weber, T. N. Rescigno, et al., Phys. Rev. A, 81(2010)011402R.

3 - 8 Unambiguous Detection of Interatomic Coulombic Decay of Ne Dimer*

Yan Shuncheng, Zhang Pengju, Shen Lili and Ma Xinwen

A famous energy transfer mechanism is termed Interatomic Coulombic Decay (ICD)^[1,2], in which an initial vacancy is firstly created in an inner shell of atom A, then the excess energy from the drop of an outer shell electron into the vacancy can be transferred to the neighboring atom B via virtual photons.

In contrast with numerous X ray and heavy ion experiments, the ICD investigation by electron impact is very scarce. And the corresponding evidence is only obtained based on electron energy spectrum subtraction^[3]. Now, in order to get the ICD evidence more directly, we performed the fragmentation experiment of Ne dimers by 380 eV electron impact at the Reaction Microscope in the Institute of Modern Physics, CAS^[4].

By detecting the Ne^+/Ne^+ ions and the emitted electron in coincidence, the momenta and energy of all charged products, as well as the Kinetic Energy Release (KER), are obtained. As shown in Fig. 1, the relationship of the

* Foundation item: National Natural Science Foundation of China (11404325, 11304325, U1532129)