

proton partial widths for those states above the proton threshold in  $^{18}\text{Ne}$ .




Fig. 2 Identification plot for the recoiled light particles at  $\theta_{\text{lab}} \approx 10^\circ$ .

measured by using three sets of  $\Delta E$ - $E$  Si telescopes at averaged angles of  $\theta_{\text{lab}} \approx 3^\circ$ ,  $10^\circ$  and  $18^\circ$ , respectively. Fig. 2 shows the particle identification by using the  $\Delta E$ - $E$  method and the recoiled protons are clearly identified. The energy calibration for the silicon detectors was performed by using a standard triple  $\alpha$  source. Due to the pulse height defect, a secondary proton calibration based on the  $\alpha$  calibration was made using proton beams of various energies. In addition, at secondary target position, an Ar gas target at 158 mbar was used in a separate run for evaluating the background contribution. Nowadays, the data analysis is still in progress.

## References

- [1] W. H. G. Lewin, J. V. Paradijs, R. E. Taam, *Space Science Reviews*, 62(1993)223.
- [2] R. K. Wallace, S. E. Woosley, *Astrophys. J. Suppl.*, 45(1981)389.
- [3] K. I. Hahn, et al., *Phys. Rev.*, C54(1996)1999.
- [4] B. Harss, et al., *Phys. Rev.*, C65(2002)035803.
- [5] Y. Yanagisawa, et al., *Nucl. Instr. and Meth.*, A539(2005)74.

## 2 - 21 Progress Report on Lanzhou Penning Trap (LPT)

Huang Wenxue, Tian Yulin, Wang Junying, Wang Yongsheng, Sun Yuliang, Zhao Jianmin  
Wang Yue, Wu Wei, He Yuan, Ma Lizhen, Xu Hushan and Xiao Guoqing

The Lanzhou Penning Trap (LPT) is an ion trap facility that is presently under construction at the Institute of Modern Physics (IMP). Its main goal is to perform direct mass measurements on fusion-evaporation residues and if possible for the heavy isotopes. Mass measurements of stable and exotic nuclei allow determination of nuclear binding energies and hence provide important data for nuclear physics and astrophysics. The following progresses on the LPT have been achieved in 2012:

(1) All parts of the whole LPT beam line have been assembled and put together, including two Penning traps, beam transport devices, and other devices for vacuum pumping, support, and so on. Fig. 1 shows a photo of the core part.

(2) The beam line alignment has been conducted and the test results have shown that the overall accuracy is within  $\pm 0.5$  mm and mostly  $\pm 0.2$  mm. The vacuum tests have been taken and an ultra high vacuum  $\sim 10^{-6}$  Pa has been achieved. The gas feeding system to the purification trap has been also tested and a pressure ratio of  $\sim 35$  has been measured between the purification trap and the measurement trap. At last the feedthrough for the electrodes has been finished and assured by using a 500 V tramegger.

(3) The manufacturing and winding of the home-made 7 T superconducting magnet have been finished and the coils have been tested by liquid nitrogen.




Fig. 1 Photo of the two Penning traps.

## 2 - 22 Isospin Effect on Transverse Flow for Isobaric Fragments from Nuclear Collisions at Intermediate Energy

Liu Xingquan, Wada Roy, Chen Zhiqiang, Huang Meirong, Lin Weiping  
Zhang Suyalatu, Jin Zengxue, Han Rui, Liu Jianli and Shi Fudong

Isospin effects, especially from system isospin asymmetry and fragment isospin asymmetry, are introduced into transverse flow extraction mainly via various physical quantities such as symmetry potential, NN cross section, and coulomb force. Since lots of efforts have been carried out to study the isospin effect by comparing the colliding pairs with different isospin in both experiment<sup>[1-5]</sup> and theoretical calculations<sup>[6-8]</sup>, in these paper, we focus on the fragment isospin effect on the flow for isobaric fragments.




Fig. 1 Transverse flow in the reaction  $^{40}\text{Ca} + ^{40}\text{Ca}$  at 35 A MeV extracted from different CoMD calculations versus A number. Isobaric fragments with  $I(N-Z) = -1-2$  are displayed with  $\triangle$   $I = -1$ ,  $\circ$   $I = 0$ ,  $\blacktriangle$   $I = 1$ ,  $\bullet$   $I = 2$ , respectively.